

the exception of the Heterohelicidae, which have had their own separate evolutionary development. *Globocornusa daubjergensis* (Brönnimann) is classified with the Guembelitrinae in this paper.

Four lineages within the genus *Globorotalia* are differentiated: one leading toward low-conical, keeled and non-keeled forms (*G. spiralis-pusilla-convexa-broedermanni*) became extinct during the middle Eocene; a second leading toward compressed, keeled, and non-keeled forms (*G. compressa-ehrenbergi-pseudomenardii-chapmani-planonica-pseudoscitula*) became extinct at the end of the middle Eocene; a third is postulated as having led from low- to high-conical forms ("conical globorotaliids") which became extinct at the end of the middle Eocene; and a fourth lineage is postulated as having led to the Neogene and Recent globorotaliid faunas.

The acarininids appear to be characterized by species which are distinguished from *Subbotina* and *Globorotalia* by a strongly spinose wall texture. Two, and possibly three, branches within this lineage appear to lead independently, through parallel trends, to the development of the polytypic, but monophyletic, genus *Truncorotaloides*. In terms of normal criteria of divergence (gaps) and monophyly, there seems to be justification for recognition of the acarininids as a genus, provided the genus were emended to include primarily spinose forms. The genus *Truncorotaloides* also is accepted as valid, although it seems unlikely that it warrants a separate subfamily.

BERNER, ROBERT A., Yale University, Department of Geology, New Haven, Connecticut

CHEMICAL DIAGENESIS OF SOME MODERN CARBONATE SEDIMENTS

Mineralogical determinations and pore-water chemical analyses have been conducted on fine-grained carbonate sediments from Bermuda and southern Florida. For the pore waters of normal marine sediments, the results clearly demonstrate no change within experimental error of the ratios Mg^{++}/Cl^{-} and Sr^{++}/Cl^{-} from values found in the overlying sea water. Also, there is a lack of consistent change in carbonate mineralogy with depth. No evidence was found for the recrystallization of metastable aragonite or high-Mg calcite to low-Mg calcite and dolomite in sediments which have been in contact with sea water since deposition. Carbonate saturo-meter measurements indicate equilibrium of the pore waters with low-Mg calcite, even though it is a minor phase in many samples. This conclusion is corroborated by calcium, alkalinity, and pH determinations, and by laboratory studies of the solution behavior of the sediments. Values of dissolved Ca^{++} lower than those expected for simple $CaCO_3$ solution by dissolved CO_2 , in sediments rich in H_2S , can be explained partly by the addition of excess HCO_3^{-} from bacterial sulfate reduction and partly by the precipitation of calcium phosphate.

Pore waters from the brackish mangrove-swamp sediments of the Everglades contain a slight excess of dissolved Mg^{++} over that expected from the Mg^{++}/Cl^{-} ratio of the overlying brackish waters and of sea water. This suggests that transformation of high-Mg calcite to low-Mg calcite takes place in these sediments by interaction with fresh water. Loss of Mg from calcite in fresh water is directly demonstrated by the chemistry of Bermuda cave waters.

The lack of diagenetic transformation of carbonate minerals in sea water may be caused mainly by the interaction of dissolved Mg^{++} with the surfaces of the mineral grains.

BRYANT, WILLIAM R., Texas A&M University, College Station, Texas, and HARDING, JAMES L., Nichols State College, Thibodaux, Louisiana

GEOTECHNICAL PROPERTIES OF MARINE SEDIMENTS FROM GULF OF MEXICO

Sediment cores collected from three areas of the Gulf of Mexico—the Louisiana continental shelf, the Mexican shelf-slope, and the abyssal plain—were examined to determine their geotechnical properties.

As these three areas have diverse sedimentation rates ranging from rapid in proximity to the Mississippi delta to very slow on the abyssal plain, the cores were studied with the expectation that a correlation would be evident between rate of deposition and the following properties: shear strength, wet density, water content, grain-size distribution, and porosity.

The cores examined from the Louisiana and Mexican shelves and the abyssal plain exhibited similar characteristics, with shear strength in the upper 10 feet of the cores averaging about 90 lbs./sq. ft.

The shear-strength values of the cores taken on the Mexican slope average much higher (170 lbs./sq. ft.), with some zones having a shear strength as high as 800 lbs./sq. ft. Such a high shear strength is caused by authigenic cementation and the presence of volcanic shards.

Statistically speaking, the remaining parameters measured showed less correlation between zones than did shear strength.

It is assumed from this study that, although the shelf and abyssal zones are being subjected to different sedimentation rates, the processes of sedimentation are such that sediments on the Louisiana and Mexican shelves and the abyssal plain are underconsolidated as opposed to the slope sediments, which are in a normal to overconsolidated state.

BUCHANAN, GEORGE S., Husky Oil Company, Houston, Texas, and SCHWAB, ROLAND F., Western Tunisia Company, Tunis, Tunisia

MIDDLE JURASSIC TROUGH OF CENTRAL TUNISIA

Upper Jurassic rocks are widespread in the surface outcrops of Tunisia and adjacent areas in Algeria and Libya. Deep drilling in central Tunisia during the past 15 years has revealed the presence of a thick rock sequence of limestone and calcareous shale which are recognized as Middle Jurassic (Callovian, Bathonian, and Bajocian), and which, in some places, are 1,500 meters thick.

To facilitate better understanding and appreciation of the geological framework and setting for the depositional trough which accommodated this thick series of sediments, maps showing the areas of Triassic outcrops and the thickness of these beds at the surface and in the subsurface are presented. The environment and nature of Triassic sediments are presented, followed by a discussion of the entire Jurassic System in Tunisia. Emphasis is placed on the Middle Jurassic and on the possible role of "Dogger" beds of early Middle Jurassic age serving as reservoir or source beds for oil and gas.