

quartz, as well as small intraclasts or oölites, trapped between laminations and in small cut-and-fill structures.

In the mats, laminae are essentially smooth and parallel. Mats are composed almost entirely of fine silt-size dolomite and commonly bear desiccation features which suggest a calm shallow environment. This is in contrast to the presence of intraclasts and algal clasts in the mounds which indicate a relatively high-energy environment. Together, both forms of stromatolites suggest a shallow, intertidal, locally hypersaline environment very much like that of the areas where algal stromatolites occur today.

DENNISON, JOHN M., University of Tennessee, Knoxville, Tennessee, and MORRISON, ANNE K., Hamburg, Germany

FACIES RELATIONS OF HYSTRICHOSPHERES IN ONESQUETHAW STAGE (DEVONIAN) OF CENTRAL APPALACHIANS

Eighteen stratigraphic sections in the Valley and Ridge area of Virginia (10), West Virginia (6), Maryland (1), and southern Pennsylvania (1) were sampled in detail to establish the distribution of hystrichospheres in the Onesquethaw Stage of that region. The localities were chosen carefully relative to the stratigraphic and paleogeographic interpretation of Dennison (1961). Rock-stratigraphic units sampled include the Tioga Metabentonite, Huntersville Chert (including Bobs Ridge Sandstone Member), Onondaga Limestone, and Needmore Shale (with three subfacies: calcitic shale and limestone, calcitic shale, and Beaver Dam black shale).

No hystrichospheres were observed in the Bobs Ridge or other glauconitic sandstone beds in the Huntersville Chert. Only one fragmentary form was seen in the Tioga Metabentonite. The Onondaga Limestone, the three subfacies of Needmore Shale, and the chert of the Huntersville all contain hystrichospheres. Hystrichospheres are rare in Huntersville Chert adjacent to the Monroe Island of middle Onesquethaw age but are abundant in chert elsewhere. Similarly, they appear to be absent in all three subfacies of Needmore Shale adjacent to Monroe Island (at Covington, Virginia). Farther from land, in a basin toward the east between the island and a source of clastic mud near Baltimore, hystrichospheres are abundant in the calcitic shale and common in black-shale, calcitic-shale, and limestone subfacies.

Nineteen species have been identified, assignable to *Hystrichosphaeridium*, *Veryhachium*, *Polyedryxium*, *Michrystridium*, and *Cymatiosphaera*. *Michrystridium* seems more abundant and *Veryhachium* less common in the calcitic-shale and limestone subfacies of the Needmore Shale than in the other hystrichosphaerid-bearing rocks. *Cymatiosphaera* is less abundant in Beaver Dam black shale than in other rock types. No other statistically valid distribution trends could be observed.

There is no evidence that hystrichospheres can be used to subdivide the Onesquethaw Stage in this region.

DONATH, FRED A., Columbia University, New York, New York

EFFECTS OF DUCTILITY AND PLANAR ANISOTROPY IN FOLDING OF ROCK LAYERS

The mechanisms of folding that can operate within

a sequence of layered rocks are dependent on the relative ductilities and on the nature of inherent anisotropy in the rocks. Flexural mechanisms require the presence of mechanical anisotropy, and folding is effected by slip between layers, by flow within layers, or by a combination of the two. Flexural folding represents a true bending of layers. Passive mechanisms operate only in rock sequences that are mechanically isotropic. This condition may exist either because of the absence of effective planar anisotropy or because of the ineffectiveness of anisotropy resulting from high ductility. The geometry of flow in passive folding reflects only the stress field and velocity gradients existing during deformation. Flow occurs across the layering, and the layering serves only to record the relative displacements. A gradational mechanism, by which certain layers are flexed in response to passive behavior in the associated rocks, causes quasi-flexural folding. This mechanism operates primarily in layered sequences characterized by high-ductility contrast, and produces strongly disharmonic folding.

Because changing environmental conditions can alter both the ductilities of the rocks involved and the effectiveness of planar anisotropy, the mechanism that initiates folding may be superceded by one or more other mechanisms during the course of deformation. The effect of layering in the folding process decreases with increasing ductility, and every gradation exists between ideally flexural and ideally passive folding.

DOTT, ROBERT H., JR., University of Wisconsin, Madison, Wisconsin

COHESION AND FLOW PHENOMENA IN CLASTIC INTRUSIONS

Physical differences between water-saturated cohesionless sand and cohesive fine mud account for many sedimentary structures. Relative bulk cohesion is controlled by relations of intergranular adhesion caused by electromagnetic or Van der Waal's forces versus the disturbing effect of forcible migration of pore fluid or gas. Where expulsion of fluid is slow, mud retains greater cohesion than sand. If sand is loosely packed, movement of fluid, especially if sudden, may cause spontaneous liquefaction and conversion to a viscous slurry with negligible strength.

Empirical evidence of major differences of cohesion comes from abundance of shale pebbles, load structures, current sole marks, contorted stratification, and clastic intrusions. These are most characteristic of alternating sand-mud sequences which possessed many inherent-strength discontinuities. All degrees of liquefaction, flow, intrusion, and stopping are "frozen" in rocks—from local in-place liquefaction of sand strata to immense dike complexes and sand volcanoes. Clastic dikes are of special interest. Typically they contain fine sand, but in a few places may have coarse gravel. Wall rock is generally mud, and stoped cohesive mud xenoliths are common; some dikes split and even rejoin along strike. Internal lamination and some grain alignment may occur, and these suggest chiefly laminar flow of viscous suspensions. A few dikes display excellent groove and flute marks on their sides, evidencing scouring of cohesive wall rock during intrusion and indicating direction of intrusion. Some large dikes completely evacuated their source stratum with concomitant subsidence of all overburden.

Important relative age criteria are provided. Most dikes are straight, suggesting that compaction was completed largely before intrusion. In fact, compac-